



NEW DETRITAL ZIRCON U-PB AGES ON NEOPROTEROZOIC TANDILIA SEQUENCES, RÍO DE LA PLATA CRATON, ARGENTINA

VII SSAGI
South American Symposium
on Isotope Geology
Brasília, 25th - 28th July 2010

Carlos A. Cingolani^{1,2}, Norberto J. Uriz², Farid Chemale Jr.³

¹ Centro de Investigaciones Geológicas (UNLP-CONICET), La Plata, Argentina. coringolani@mdp.unlp.edu.ar / carloscingolani@yahoo.com
² División Científica de Geología Museo de La Plata, Paseo del Bosque, 1900-La Plata, Argentina.
³ Núcleo de Geología, Universidade Federal do Sergipe, Brazil.



University of La Plata



Figure 1: Southeastern South American sector showing the inferred limits of the Rio de la Plata craton region (Cordani, 2009) and the position of the Tandilia System.

INTRODUCTION

The geological evolution of Tandilia comprises mainly a juvenile igneous-metamorphic Paleoproterozoic basement named Buenos Aires Complex, which is covered by thin Neoproterozoic to Early Paleozoic sedimentary units which displays sub horizontal bedding (Fig. 1). This work is related to introduce the new detrital zircon U-Pb ages from two siliciclastic Neoproterozoic sequences called Villa Mónica and Cerro Negro Formations from the Barker region, and offer the comparison provenance analysis of these data with previous contributions in a paleogeographic interpretation.

ANALYTICAL METHODS

For isotopic dating, all zircon grains were mounted in 2.5 cm-diameter circular epoxy mounts and polished down until the zircons were just revealed. Images of zircons were obtained using the optical microscope (Leica MZ 125) and backscatter electron microscope (JEOL JSM 5800). Zircons were dated with a LA microprobe (New Wave UP123) coupled to a MC-ICP-MS (Neptune) at the Laboratorio de Geología Isotópica, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil.




Figure 1: Geological sketch map of the Barker region and the location of the study samples. The inset shows the relative location of the area in the south-centre of the Tandilia belt. Geology based on Leveratto and Marchese (1983)

RESULTS

The U-Pb data is shown in Fig. 2 and could be summarized as follow, taken into account that the sedimentary cover of Tandilia was characterized by four marine transgressions.

a. First marine transgression (Villa Mónica Fm): The sample analyzed here was collected at the base of the unit composed by a quartzite-type rock in the Águilas Range at the Barker region (Fig. 1). The study zircon grains have subrounded to rounded shapes suggesting an intense hydraulic transport. Sixty dated zircons yield a unimodal population with a main peak at 2.25 Ga with typical Paleoproterozoic ages between 1.97 and 2.28 Ga (90%). These data confirm the previous one by Rapela *et al.* (2007) and Gaucher *et al.* (2008) implying that the detritus are coming mainly from the Rhyacian to Orosirian sources, commonly called in South America a "Transamazonian cycle". Detrital zircons from the Archean source are present in a 10%. The youngest detrital zircon of sample 05-CA 1 yielded an age of c. 2.0 Ga.

b. Third marine transgression (Cerro Negro Fm, "phosphate member" at the base of the unit): Fifty-eight detrital zircons were analyzed from a quartz-rich sandstone sample as we can see on Fig. 2. Zircon grains are very rounded without faces preserved; fractures are common but the crystals are complete. The frequency diagram shows a polymodal distribution pattern spanning most of the Archean, the Palaeoproterozoic, and Mesoproterozoic. The Palaeoproterozoic zircons at 2.26 to 1.7 Ga is the most important population with peaks at 2.14, 2.05 and 1.97 Ga and comprising 75 % of all dated grains. These zircon grains are coming also from Rhyacian to Orosirian sources. The Mesoproterozoic zircon peaks are at 1.53, 1.21 and 1.16 Ga and span from Calymnian to Stenian ages (20 %). Three Archean peaks (less important than the previous mentioned) span from 2.56, 2.73 and 2.90 Ga (5 %). The youngest detrital zircon of sample 05-CN 3 yielded an age of c. 1.12 Ga.




Figure 2: U-Pb Concordia diagrams for detrital zircon grains of the Villa Mónica Fm (left) and Cerro Negro Fm (right) samples. The backscattered electron images of some analyzed zircons on the bottom of both Concordia diagrams show the main typology characteristics of dated grains and LA-ICP-MS spots.



DISCUSSION

As we can see on Fig. 3, the data for the first marine transgression (Villa Mónica Fm) may be compared with the data of Rapela *et al.* (2007), who report 55 detrital zircon ages for one sample from the same area. Its zircon population shows predominantly Paleoproterozoic with peak at 2.24 Ga, but it includes a subordinate Mesoproterozoic peak centred at 1135 Ma. The data published by Gaucher *et al.* (2008) are coming from a sample that was collected at the Villa Mónica Quarry which is the stratotype of this unit, about 100 km from Barker region. These authors report 81 dated zircons that yield also a unimodal population centred at 2146 Ma with typical "Transamazonian ages". Based on these coincident results it is clear that the source of the Villa Mónica Fm must have been mainly the underlying Palaeoproterozoic Buenos Aires Complex well exposed in Tandilia belt.

The data of the third marine transgression over Tandilia belt (Cerro Negro Fm) could be compared (Fig. 3) with the quartz-rich sample from the underlying unit called Cerro Largo Fm (second transgression) obtained by Gaucher *et al.* (2008), these authors report more than hundred concordant zircons yielding a complex polymodal distribution spanning the Archean, the Palaeoproterozoic, and Mesoproterozoic ages.

Palaeoproterozoic zircons represent the most important population and comprise 58% of all dated grains. These ages are roughly coincident with the orogenic events, recognized in Tandilia by Hartmann *et al.* (2002). It is important to note that the minor zircon population yielding Archean ages is present in Villa Mónica, Cerro Largo and Cerro Negro Fms. That confirms that these units contain in different percentages the oldest detrital zircons dated for the Tandilia region.

The most conspicuous ages of detrital zircons in the Neoproterozoic sedimentary units study here are Palaeoproterozoic, which match the widespread "Transamazonian orogenic cycle" during Rhyacian to Orosirian ages. The provenance of the Archean detrital zircons could be derived from the Nico Pérez terrane (Uruguay) where is known to contain extensive Palaeo- to Neoproterozoic magmatic and metamorphic rocks (Gaucher *et al.*, 2008). The abundance of Mesoproterozoic zircons in the Cerro Largo and Cerro Negro samples need to be explained in a paleogeographical detail. Another striking feature to be mentioned is the absence of Neoproterozoic zircons suggesting that this region may be a passive margin of the Pan-African-Brasiliano Cycle.




Figure 3: Comparative U-Pb provenance patterns for sedimentary samples from the Tandilia belt. The curves are relative probability trends. Data in stratigraphical relative position from base to top are based on Rapela *et al.* (2007); Gaucher *et al.* (2008), and the obtained in this work.

REFERENCES:

Dalla Saldia LH, Spalletti LA, Poiré DG, De Barrio R, Echeveste H, Benialgo A. 2006. Tandilia. In: Temas de Geología Argentina 1. Serie Correlación Geológica, 21:17-46. San Miguel de Tucumán.
Gaucher C, Finney SC, Poiré DG, Valencia VA, Grove M, Blanco G, Pamoukaghlian K, Gómez Peral L. 2008. Detrital zircon ages of Neoproterozoic sedimentary successions in Uruguay and Argentina: Insights into the geological evolution of the Rio de la Plata Craton. *Precambrian Research* 167, 150-170.
Hartmann LA, Santos JOS, Cingolani CA, McNaughton NJ. 2002. Two Paleoproterozoic Orogens in the Evolution of the Tandilia Belt, Buenos Aires, as evidenced by zircon U-Pb SHRIMP geochronology. *International Geology Review*, 44: 528-543.
Lanza HA, Hugo C. 1987. Descubrimiento de fosforitas sedimentarias en el Proterozoico superior de Tandilia, Buenos Aires, Argentina. *Revista de la Asociación Geológica Argentina* 42:3-4 y 417-428.
Leveratto MA, Marchese HG. 1983. Geología y estratigrafía de la Formación La Tinta (y homónimas) en el área clave de Sierra de La Tinta-Barker-Villa Caicue-Arroyo Calaveras, Provincia de Buenos Aires. *Revista de la Asociación Geológica Argentina* 38: 235-247.
Poiré DG, Gaucher C. 2009. Lithostratigraphy. Neoproterozoic-Cambrian evolution of the Rio de la Plata Palaeocontinent. In: Gaucher C *et al.* (eds) *Neoproterozoic-Cambrian tectonics, global change and evolution. Developments in Precambrian Geology*, 16, 4, 2, 87-101. Elsevier.
Rapela CW, Pankhurst RJ, Casquet C, Fanning CM, Baldo EG, González-Casado JM, Galindo C, Dahlquist J. 2007. The Rio de la Plata Craton and the assembly of SW Gondwana. *Earth Sci Rev* 83, 49-82.

Acknowledgments: We are grateful to CONICET (Argentina) and CNPq (Brazil) for financial support.